

Parametrizations of Quark Confinement in Production Processes

C.S. Fischer*

Institute for Theoretical Physics, University of Tübingen
Auf der Morgenstelle 14, 72070 Tübingen, Germany

Baryons are described as bound states of quarks and scalar as well as axialvector diquarks. In order to effectively parametrize confinement several *ansätze* for the dressing functions of the constituent propagators are used. The corresponding results for electromagnetic and strong form factors differ only slightly. On the other hand observables from production processes show the permissibility of different dressing functions.

The aim of the present investigation is to set up a model describing baryon properties at intermediate energies, where great experimental progress has been made in the last decade. In order to capture essential features of QCD at least two requirements should be met: the model has to be Poincaré covariant and it has to incorporate confinement to avoid unphysical thresholds. The diquark-quark model fulfills both demands.

Quark-quark correlations inside the baryon are dominated by separable contributions that carry the quantum numbers of the scalar and the axialvector diquark [1]. Therefore baryons can be described by a quark-diquark Bethe-Salpeter (BS) equation, with the constituents interacting via quark exchange [2,3]. We solve the BS equation without reduction (cf. [4]) to obtain covariant vertex functions for octet and decuplet baryons.

The central input of the model is the *ansatz* for the dressing function of the quark and diquark propagators. We employ three different methods to parametrize confinement: first we use an exponential dressing function to shift the poles of the bare propagators to timelike infinity ('exp-propagator'), second we split the real poles into complex conjugate pairs ('cc-propagator') and third we employ a nonanalytic dressing function that removes the poles completely ('tanh-propagator') [5].

The electric and magnetic form factors of the

proton as well as the mass spectrum of the octet baryons serve to fix the parameters of the model. Special care is taken of electromagnetic current conservation [6]. Although the contributions from the axialvector diquark to the vertex functions of the nucleon are at the 10% level, it turns out that they are vital to reproduce the magnetic moments [7]. A summary of the electromagnetic properties of the nucleon in our model can be found in [8].

Having fixed the parameters, the results for the strong form factors are predictions. We obtain a soft pion nucleon form factor $g_{\pi NN}$ and good agreement with the experimental value at zero momentum. Due to flavour symmetry breaking the results for the kaon nucleon lambda form factor $g_{KNA}(p^2)$ are smaller than the one for $g_{\pi NN}(p^2)$ for all momenta. In general all three *ansätze* for the dressing functions of the propagators lead to similar results for form factors [5].

This is different for production processes. In the diquark spectator approximation there are two types of diagrams that describe kaon photoproduction, $\gamma p \rightarrow \Lambda K$. In the s-channel diagram both, photon and kaon, couple to a single quark line, whereas in the t-channel the photon couples directly to the kaon. Fig. 1 shows that the parametrization with the exponential dressing function overshoots the data by orders of magnitude. This is because one of the propagators in the s-channel diagram is probed for large timelike momenta where the exponential dressing blows up very quickly. We therefore consider the

*Talk given at the workshop *The Spin Structure of the Proton*, Trento, July 23rd to 28th, 2001.




Figure 1. Total cross section of $\gamma p \rightarrow \Lambda K$ compared to the data of the SAPHIR-collaboration [10].

exponential dressing *ansatz* to be ruled out for the whole class of models that resolve baryons into its quark substructure [5]. The other two parametrizations describe the data reasonably well. Here the t-channel diagram dominates. It has been shown, however, that interference between the two channels even leads to a good description of the substructure of the data [9].

For associated strangeness production, $pp \rightarrow pK\Lambda$, we obtain a similar picture with respect to the dressing functions. Again the exponential dressing overshoots the data massively, whereas the cc-propagator and the tanh-propagator give cross sections with the right order of magnitude. The spin depolarisation tensor D_{NN} provides a measure for the relative strength of diagrams with pion and kaon exchange between the two protons. Pion exchange leads to positive values of D_{NN} whereas kaon exchange gives negative contributions. Fig. 2 shows, that the experimental data seem to favour kaon exchange, whereas pion exchange diagrams dominate in our model [5]. We regard this as a sign for missing contributions in the kaon exchange channel.

Acknowledgements

I thank S. D. Bass, A. De. Roeck and A. Deshpande for organising this very interesting




Figure 2. Depolarisation tensor D_{NN} for $pp \rightarrow pK\Lambda$ compared to the data of the DISTO-collaboration [11].

workshop and for the invitation. I am grateful to S. Ahlig, R. Alkofer, M. Oettel, H. Reinhardt and H. Weigel for their collaboration on the work presented here. Thanks for the support by the European Graduate School Basel-Tübingen and by COSY under contract no. 4137660.

REFERENCES

1. I. Wetzorke and F. Karsch, hep-lat/0008008.
2. M. Oettel et al., Phys. Rev. **C58** (1998) 2459.
3. R. Alkofer and L. von Smekal, Phys. Rep. **353** (2001) 281; C. D. Roberts and S. M. Schmidt, Prog. Part. Nucl. Phys. **45S1** (2000) 1.
4. M. Oettel and R. Alkofer, Phys. Lett. **B484** (2000) 243.
5. S. Ahlig *et al.*, Phys. Rev. D **64** (2001) 014004.
6. M. Oettel, M. Pichowsky and L. von Smekal, Eur. Phys. J. **A8** (2000) 251.
7. M. Oettel, R. Alkofer and L. von Smekal, Eur. Phys. J. **A8** (2000) 553.
8. R. Alkofer and M. Oettel, hep-ph/0105320.
9. S. Ahlig *et al.*, Prog. Part. Nucl. Phys. **44** (2000) 361.
10. M. Q. Tran *et al.* [SAPHIR Collaboration], Phys. Lett. **B445** (1998) 20.

11. F. Balestra *et al.* [DISTO Collaboration],
Phys. Rev. Lett. **83** (1999) 1534.